

AD-A157 332

MOLECULAR MECHANISMS INVOLVED IN TISSUE SWELLING DUE TO 1/1
INJURY AND DUE TO (U) PENNSYLVANIA HOSPITAL
PHILADELPHIA DEPT OF MOLECULAR BIOLOGY.. G N LING


UNCLASSIFIED

28 MAY 85

F/G 6/16

NL





MICROCOPY RESOLUTION TEST CHART
 NATIONAL BUREAU OF STANDARDS-1963-A

AD-A157 332

OFFICE OF NAVAL RESEARCH

Contract N00014-79-C-0126

Task No. NR 207-017

FINAL REPORT

Molecular Mechanisms Involved in Tissue Swelling Due to Injury and Due to Exposure to Low Temperature and Massive Water and Electrolyte Loss in Diarrheal Disorders

by

Gilbert N. Ling, Ph.D.

Department of Molecular Biology
Pennsylvania Hospital
Eighth and Spruce Streets
Philadelphia, PA 19107

DTIC
ELECTE
JUL 17 1985
S D
G

*Distribution of this report is unlimited

DTIC FILE COPY

45 10 36 054

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO. <i>AD-A157332</i>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Molecular Mechanisms Involved in Tissue Swelling Due to Injury and Due to Exposure to Low Temperature and Massive Water and Elec- trolyte Loss in Diarrheal Disorders		5. TYPE OF REPORT & PERIOD COVERED Final 12-1-79 - 4-1-85
7. AUTHOR(s) Gilbert N. Ling, Ph.D.		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Molecular Biology, Pennsylvania Hos- pital, Eighth & Spruce Streets Philadelphia, PA 19107		8. CONTRACT OR GRANT NUMBER(s) N00014-79-C-0126
11. CONTROLLING OFFICE NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 207-017 Distribution of this report is unlimited
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE May 28, 1985
		13. NUMBER OF PAGES 9
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Unlimited		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) ion adsorption, water polarization, ATP, cell injury, NMR		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The ONR-supported work conducted in this laboratory has helped to esta- blish a major change in our fundamental concept of the living cell. This basic change would affect virtually every phase of biomedical research. Not surprising, this new development has already produced a major practical advance (NMR imaging) of benefit to military as well as civilian populations.		

a. Summary of all research accomplished

The summary only refers to work accomplished since 1979. Work accomplished before 1979 has been repeatedly presented in the individual reports and is also summarized in the monograph published in 1984 (see a.1.).

a.1. Publication of the monograph, In Search of the Physical Basis of Life (791 pages), Plenum Publishing Corporation. In this the essence of the association-induction hypothesis (a general theory of the living cells presented by the principal investigator in 1962) is presented along with results of thirty years of experimental testing in the historical perspective of mans' search for the understanding of the physical basis of life. A copy of this volume has been submitted to the Office of Naval Research in the care of Dr. J. Majde.

a.2. The following are brief summaries of specific research accomplished, largely published in the publications listed below under c.

a.2.1. Model Studies

a.2.1.1. Proteins: Alteration of the physical state of water in vitro by proteins when they exist in the fully extended state for structural reasons or in response to the presence of denaturants

In the altered physical state, the bulk phase water has reduced solubilities for solutes normally excluded from living cells (e.g., Na^+ , sucrose, glycine) but normal solubility for solutes not excluded by living cells (e.g., urea). When the protein chains are locked in α -helix or β -structure with the water polarizing NHCO groups marked, the physical state of water is essentially normal.

a.2.1.2. Polymers

Solute exclusion properties demonstrated in fully extended protein chains were also demonstrated in synthetic polymers which possess the basic attributes required by theory (possession of fixed charged groups (CO, NH) at proper distances apart).

a.2.1.3. Polymers

Water which is dominated by polymers and has reduced solubilities for Na^+ , etc., showed moderately longer NMR rotational correlation time (τ) approximately equal T_1 and T_2 . This finding is of crucial importance toward our attempt to interpret the T_1 and T_2 of water protons in living cells.

a.2.1.4. Proteins and Polymers

Proteins with fully extended polypeptide chains and polymers which exclude Na^+ , etc., also demonstrated unusually high osmotic activity in confirmation of theory.

a.2.1.5. Volume regulation of living cells and models of living cells

Cell volume is controlled by concentrations of Na salts in the external solution even though there is no intact cell membrane. This behavior (i.e., swelling and shrinkage without a classic "semipermeable membrane") is confirmed in studies of solutes of polymers which partially exclude Na salts.

a.2.1.6. Freezing and thawing pattern of water dominated by polymer which reduces water solvency for Na^+ , etc.

It was demonstrated that native proteins (up to 50% (w/v)) which has little effect on water solvency for Na^+ , also exercises very little effect on the freezing and thawing points of water. Freezing and thawing points are progressively reduced by proteins or polymers that reduce water solvency for Na^+ , etc. This finding is also of historical importance because it gives new significance to freezing point depression in water of living cells done in the past and so far wrongly interpreted.

a.2.1.7. β - and γ -carboxyl groups

β - and γ -carboxyl groups carried by proteins stoichiometrically and preferentially bind alkali metal ions when these anionic groups are "unmasked" after neutralization of competing fixed cations (e.g., ϵ -amino groups).

This finding was also of historic importance because it has traditionally been believed that proteins bind little K^+ , Na^+ , or other alkali-metal ions and binding of K^+ to β - and γ -carboxyl groups is a main postulate of the association-induction hypothesis.

a.2.2. Living Cells

a.2.2.1. Confirmation of the "Universality Rule"

"If the solubility of one normally partially excluded solute in a living cell goes up or down, the solubilities of all other normally excluded solutes will change *pari passu*". This rule based on the AI hypothesis has been confirmed in frog muscle very slowly dying in the presence of a low concentration of metabolic poison (0.2 mM iodoacetate).

a.2.2.2. Membrane-pumps vs. cytoplasmic proteins

Red blood cell vesicles with intact membrane of healthy Na, K activated ATPase but little or no cytoplasm ("white ghosts") do not reaccumulate K^+ and extrude Na^+ in the presence of ATP. Only "red ghosts" which have retained hemoglobin and other cytoplasmic proteins reaccumulate K^+ and extrude Na^+ . Furthermore the extent of K^+ reaccumulated and Na^+ extruded are quantitatively dependent on the cytoplasmic proteins present.

a.2.2.3. Water in cancer cells is less able to exclude pentose probes while most living cells do.

a.2.2.4. Demonstration that the NMR relaxation times are partly due to paramagnetic ions present in the cells.

a.2.2.5. Longer NMR relaxation times T_1 and T_2 in cancer cells largely reflect a lower paramagnetic ion content.

- a.2.2.6. Swelling of tissues suffering from anoxia and cold injury depend on the Na concentration in the medium in a qualitative manner. In the presence of fixed concentration of external Na^+ the degree of swelling increases with decrease of cell ATP content. Both sets of results confirm prediction of the AI hypothesis

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A/	



b./c. Technical reports/publications issued under contract (12-1-79/4-1-85)

1. Ling, G. N., "The Polarized Multilayer Theory of Cell Water and Other Facets of the Association-Induction Hypothesis Concerning the Distribution of Ions and Other Solutes in Living Cells" in The Aqueous Cytoplasm, pp. 23-60, ed., A. D. Keith, Marcel Dekker, Inc., NY, 1979.
2. Ling, G. N., "The Polarized Multilayer Theory of Cell Water According to the Association-Induction Hypothesis" in Cell-Associated Water, pp. 261-270, eds., W. Drost-Hanson and J. Clegg, Academic Press, Inc., NY, 1979.
3. Ling, G.N., "The Equations for Cellular Resting Potentials According to the Surface Adsorption Theory, A Corollary of the Association-Induction Hypothesis", Physiol. Chem. Phys. 11:59-64 (1979).
4. Ling, G. N., "Ion and Water Transport: Experimental Design Defended", Trends in Biochemical Sciences 4:N134-N135 (1979).
5. Ling, G. N., Walton, C., and Ling, M. R., " Mg^{++} and K^+ Distribution in Frog Muscle and Egg: A Disproof of the Donnan Theory of Membrane Equilibrium Applied to the Living Cells", J. Cell. Physiol. 101:261-278 (1979).
6. Ling, G. N., and Negendank, W., "Do Isolated Membranes and Purified Vesicles Pump Sodium? A Critical Review and Reinterpretation", Persps. in Biol. & Med. 23:215-239 (1980).
7. Ling, G. N., Ochsenfeld, M. M., Walton, C., and Bersinger, T. J., "Mechanism of Solute Exclusion from Cells: The Role of Protein-Water Interaction", Physiol. Chem. Physics 12:3-10 (1980).
8. Ling, G. N., Walton, C., and Bersinger, T. J., "Reduced Solubility of Polymer-Oriented Water for Sodium Salts, Amino Acids, and Other Solutes Normally Maintained at Low Levels in Living Cells", Physiol. Chem. Phys. 12:111-138 (1980).
9. Ling, G. N., "Underestimation of Na Permeability in Muscle Cells: Implications for the Theory of Cell Potential and for Energy Requirement of the Na Pump", Physiol. Chem. Phys. 12:215-232 (1980).
10. Ling, G. N., "The Theory of the Allosteric Control of Cooperative Adsorption and Conformation Changes: A Molecular Model for Physiological Activities According to the Association-Induction Hypothesis" in Cooperative Phenomena in Biology, ed., G. Karreman, Pergamon Press, NY, pgs. 39-69, 1980.
11. Ling, G. N., and Tucker, M., "Nuclear Magnetic Resonance Relaxation and Water Contents in Normal Mouse and Rat Tissues and in Cancer Cells", J. Nat. Cancer Inst. 64:1199-1207 (1980).
12. Ling, G. N., "The Role of Multilayer Polarization of Cell Water in the Swelling and Shrinkage of Living Cells", Physiol. Chem. Phys. 12:383-384 (1980).
13. Ling, G. N., Walton, C. L., and Ochsenfeld, M. M., "A Unitary Cause for the Exclusion of Na^+ and Other Solutes from Living Cells, Suggested by Effluxes of Na^+ , D-Arabinose, and Sucrose from Normal, Dying, and Dead Muscles", J. of Cell. Physiol. 106:385-398 (1981).

14. Ling, G. N., "Water and the Living Cell as Seen from the Viewpoint of a New Paradigm" in International Cell Biology 1980-1981, ed. H. G. Schweiger, Springer-Verlag, Berlin, pps. 904-914 (1981).
15. Ling, G. N., "Oxidative Phosphorylation and Mitochondrial Physiology: A Critical Review of Chemiosmotic Theory, and Reinterpretation by the Association-Induction Hypothesis", Physiol. Chem. Phys. 13:29-96 (1981).
16. Ling, G. N., "Elektrische Potentiale lebender Zellen in Die Zelle: Struktur und Funktion, 3rd ed., ed., H. Metzner, Wissenschaftlich Verlagsgesellschaft mbH, pp. 356-389, Stuttgart, Germany, 1981.
17. Ling, G. N., "Active Solute Transport Across Frog Skin and Epithelial Cell Systems According to the Association-Induction Hypothesis", Physiol. Chem. Phys. 13:356-384 (1981).
18. Ling, G. N., "Synchronous Control of Metabolic Activity by K^+ Transiently and Reversibly Liberated from Adsorption Sites During Muscle Contraction: An Extension of Association-Induction Theory", Physiol. Chem. Phys. 13:565-566 (1981).
19. Ling, G. N., and Tucker, M., "NMR Relaxation and Water Contents in Normal Tissues and Cancer Cells" in NMR in Medicine, eds. P. Diehl, E. Fluck, R. Kosfeld, Springer-Verlag, Heidelberg, pps. 161-174, 1981.
20. Ling, G. N., "The Cellular Resting and Action Potentials: Interpretation Based on the Association-Induction Hypothesis", Physiol. Chem. Phys. 14:47-96 (1982).
21. Ling, G. N., and Murphy, R. C., "NMR Relaxation of Water Protons Under the Influence of Proteins and Other Linear Polymers", Physiol. Chem. Phys. 14:209-211 (1982).
22. Ling, G. N., and Murphy, R. C., "Apparent Similarity in Protein Compositions of Maximally Deviated Cancer Cells" (Preliminary Note), Physiol. Chem. Phys. 14:213 (1982).
23. Ling, G. N., "The Association-Induction Hypothesis. A theoretical Foundation Provided for the Possible Beneficial Effects of a Low Na, High K Diet and Other Similar Regimens in the Treatment of Patients Suffering from Debilitating Illnesses", Agressologie 24:293-302 (1983).
24. Ling, G. N., "The Molecular Mechanisms of Cellular Potentials" in Structure and Function in Excitable Cells, eds. D. C. Chang, I. Tasaki, W. J. Adelman, Jr., and H. R. Leuchtag, Plenum Publishing Corp., NY, pps. 365-382, 1983.
25. Ling, G. N., and Ochsenfeld, M. M., "Studies on the Physical State of Water in Living Cells and Model Systems. I. The Quantitative Relationship between the Concentration of Gelatin and Certain Oxygen-Containing Polymers and Their Influence Upon the Solubility of Water for Na^+ Salts, Physiol. Chem. Phys. and Med. NMR 15:127-136 (1983).
26. Ling, G. N., and Murphy, R. C., "Studies on the Physical State of Water in Living Cells and Model Systems. II. NMR Relaxation Times of Water Protons in Aqueous Solutions of Gelatin and Oxygen-containing Polymers which Reduce the Solvency of Water for Na^+ , Sugars, and Free Amino Acids", Physiol. Chem. Phys. and Med. NMR 15:137-154 (1983).

27. Ling, G. N., "Studies on the Physical State of Water in Living Cells and Model Systems. III. The High Osmotic Activities of Aqueous Solutions of Gelatin, Polyvinylpyrrolidone and Poly (Ethylene Oxide) and Their Relation to the Reduced Solubility for Na^+ , Sugars, and Free Amino Acids", Physiol. Chem. Phys. and Med. NMR 15:155-165 (1983).
28. Ling, G. N., and Kwon, Young "Cold Injury-Induced Swelling of Brain and Other Tissues: Its Molecular Mechanism", Physiol. Chem. Phys. and Med. NMR 15:239-250 (1983).
29. Ling, G. N., and Zhang, Z. L., "Evidence that Na^+ in a Sulfonate Ion Exchange Resin Exists in an Adsorbed State. Its Significance for the Interpretation of NMR Data in Resins and Cells", Physiol. Chem. Phys. and Med. NMR 15:251-258 (1983).
30. Ling, G. N., and Tucker, M., "Only Solid Red Blood Cell Ghosts Transport K^+ and Na^+ Against Concentration Gradients: Hollow Intact Ghosts with K^+ - Na^+ Activated ATPase Do Not", Physiol. Chem. Phys. and Med. NMR 15:311-317 (1983).
31. Ling, G. N., and Fisher, A., "Cooperative Interaction Among Cell Surface Sites: Evidence in Support of the Surface Theory of Cellular Electrical Potentials", Physiol. Chem. Phys. and Med. NMR 15:369-378 (1983).
32. Ling, G. N., Walton, C. L., and Ochsenfeld, M. M., "Indifference of the Resting Potential of Frog Muscle Cells to External Mg^{++} in the Face of High Mg^{++} Permeability", Physiol. Chem. Phys. and Med. NMR 15:379-390 (1983).
33. Ling, G. N., and Zhang, Z. L., "Studies on the Physical State of Water in Living Cells and Model Systems. IV. Freezing and Thawing Point Depression of Water by Gelatin, Oxygen-containing Polymers and Urea-denatured Proteins", Physiol. Chem. Phys. and Med. NMR 15:391-406 (1983).
34. Zhang, Z. L., and Ling, G. N., "Studies on the Physical State of Water in Living Cells and Model Systems. V. The Warming Exothermic Reaction of Frozen Aqueous Solution of Polyvinylpyrrolidone, Poly (Ethylene Oxide), and Urea-dominated Proteins", Physiol. Chem. Phys. and Med. NMR 15:407-416 (1983).
35. Ling, G. N., "Evidence for a Significant Role of Paramagnetic Ions in the Observed NMR Relaxation Rates of Living Tissues", Physiol. Chem. Phys. and Med. NMR 15:505-510 (1983).
36. Ling, G. N., "NMR Relaxation Times of Water Protons in Normal Tissues and Cancer Cells: The Role of Paramagnetic Ions" (Preliminary Note) Physiol. Chem. Phys. and Med. NMR 15:511-513 (1983).
37. Ling, G. N., and Zhang, Z. L., "A Study of Selective Adsorption of Na^+ and Other Alkali-Metal Ions on Isolated Proteins: A Test of the Salt-Linkage Hypothesis", Physiol. Chem. Phys. and Med. NMR 16:221-235 (1984).
38. Ling, G. N., "Counterarguments Against Alleged Proof of the Na-K Pump In Studies of K^+ and Na^+ Distributions in Amphibian Eggs", Physiol. Chem. Phys. and Med. NMR 16:293-305 (1984).
39. Ling, G. N., Zodda, D., and Sellers, M., "Quantitative Relationships Between the Concentration of Proteins and the Concentration of K^+ and Na^+ in Red Cell

Ghosts", Physiol. Chem. Phys. and Med. NMR 16:381-392 (1984).

40. Ling, G. N., Baxter, J. D., and Leitman, M. I., "Effects of Adrenaline, Calcium, and Ouabain on the Resting Potential of Frog Muscle: Interpretation Based on the Theory of Allosteric Control of Cooperative Interactions Among Surface Anionic Sites", Physiol. Chem. Phys. and Med. NMR 16:405-423 (1984).
41. Ling, G. N., "Warming-Induced Hyperpolarization of Cardiac Muscle Cells and Snail Neurones: Interpretation Based on Temperature Transition of Cooperatively Linked Surface Anionic Sites Between K^+ and Na^+ Adsorbing States", Physiol. Chem. Phys. and Med. NMR 16:425-435 (1984).
42. Ling, G. N., "The Electron-Donating Strengths of Side Chains in the Determination of Protein Structure" (Preliminary Note), Physiol. Chem. Phys. and Med. NMR 16:459-460 (1984).

Monograph

Ling, G. N., In Search of the Physical Basis of Life (791 pages), Plenum Publishing Corp. New York, 1984.

d. Conclusions

The major conclusion to be drawn from the ONR-supported research is that the basic tenets of the membrane-pump theory, which has dominated biological thinking for nearly a century and is still being taught in most textbooks at all levels of education, is untenable and incorrect. The cells do not have enough energy to operate all the pumps. The ability of living cells to maintain constant levels of solutes like Na^+ , sugars, free amino acids, reflect primarily two basic mechanisms: selective adsorption on protein sites (e.g., most cell K^+) and partial exclusion from cell water at levels roughly inversely proportional to the size and complexity of the solutes and hydrated ions involved. Osmotic activity of the cell is not due to intracellular ions (which are largely adsorbed) but is due to the multilayer polarization of the cell water. Cellular resting potential is not a membrane potential with or without an electrogenic pump component; it is a surface adsorption potential. ATP adsorbed on cardinal sites on cellular proteins maintain the cellular assembly of proteins-water-ions in a high energy state - the living state. Electronic inductive effect underlies the allosteric action of ATP as well as those of Ca^{++} , hormones, and drugs that act on cells.

e. List of major accomplishments (for other accomplishments see book, cited under c)

e.1. K^+ and Na^+ distribution

(1) Provided experimental evidence that when ions are fixed in a linear chain (e.g., β - and γ -carboxyl groups), a high degree of association with counterions (e.g., K^+ , Na^+) occur. In proteins this adsorption of K^+ , Na^+ shows specificity and also cooperativity.

(2) Provided experimental proof that selective K^+ and Na^+ accumulation does not require a functional cell membrane (and postulated pumps).

(3) Provided many types of experimental studies showing what was once considered unequivocal evidence for the membrane-pump theory was wrong or equivocal.

(4) Provided proof that K^+ in frog muscle is not free. Rather K^+ is mostly adsorbed on protein sites at the A bands and Z-line in a one-site, one-ion close contact.

(5) Demonstrated that K^+ (and Na^+) distribution in frog muscle qualitatively follows the general equation for solute distribution in living cells published in 1965.

(6) Demonstrated also that K^+ , Na^+ distribution qualitatively depends on the concentration of ATP (the ultimate product of metabolism) rather than its rate of hydrolysis.

(7) Demonstrated that K^+ (and Na^+) distribution qualitatively depends on ouabain in muscle with functional membrane and without and that the pattern of distribution follows the general equation of solute distribution with ouabain binding altering the "intrinsic equilibrium constant" ($K_{\text{Na} \rightarrow \text{K}}$).

e.2. Physical state of water

(1) Introduced the polarized multilayer theory of cell water in 1964.

(2) Confirmed in frog muscle and two other model systems that the water in these systems qualitatively follows Bradley's theory of polarized multilayer adsorption.

(3) Confirmed in model systems the major postulation of the theory of cell water, i.e., (i) proteins must exist in the fully extended state with the backbone NHCO groups directly exposed to the bulk phase water and (ii) the same proteins would have little effect on the bulk phase water if the backbone NHCO are locked in intra- or intermacromolecular H bonds (see a.2.1.).

(4) Offered and confirmed new theory of cell volume control

(5) Offered a theory of cell swelling due to cold and anoxia, i.e., ATP dependent shift of the electron density of β - and γ -carboxyl groups (measured as c-value) toward increase of preference for Na^+ when compared to K^+ and fixed cations and provided experimental data supporting it.

e.3. Electrical potentials

(1) Offered a new theory of cellular electrical potential. The equation describing the potential was shown to be able to predict the experimental data which agree with the Hodgkin-Katz-Goldman equation as well as those that do not, including the "aberrant" data that led to the postulation of the electrogenic potentials.

e.4. NMR imaging

e.4.1. NMR imaging, a major advance in medical history, provides the means of non-invasive and non-injurious detection of pathological changes in human tissues. Dr. Raymond Damadian, the patent holder, wrote to me expressing the following:

"... after laboring for more than one year on the construction of our 53" superconductive magnet and on our FONAR (field focusing NMR) technique, we achieved, with great jubilation, the world's first NMR image of the live human body. The achievement originated in the modern concepts of salt and water biophysics of which you are the grand pioneer with your classic treatise, The Association-Induction Hypothesis..."

e.4.2. Our research on the molecular mechanism underlying cold- and anoxia-induced tissue swelling revealed a basic pathological change of living cells which has now been understood in terms of a theory of the living cell that does not violate basic laws of physics (as has been repeatedly shown that the conventional membrane-pump theory does). This basic knowledge provides us with indications that in treating injury-induced brain trauma and other clinical conditions the substitution of Na and chloride in the resuscitating fluid with other osmotically active solutes would be worth further investigation.

END

FILMED

9-85

DTIC